arXiv:1407.8197v1 [math.FA] 30 Jul 2014

Two-weight Norm Estimates for Multilinear Fractional Integrals in Classical Lebesgue Spaces*

Vakhtang Kokilashvili[a,b], Mieczyslaw Mastylo[c], Alexander Meskhi[a,d,1,*]

[a] *Department of Mathematical Analysis, A. Razmadze Mathematical Institute, I. Javakhishvili Tbilisi State University, 2. University Str., 0186 Tbilisi, Georgia*
[b] *International Black Sea University, 3 Agmashenebeli Ave., Tbilisi 0131, Georgia*
[c] *Faculty of Mathematics and Computer Science, Adam Mickiewicz University Poznań, and Institute of Mathematics, Polish Academy of Sciences (Poznań branch), Umultowska 87, 61-614 Poznań, Poland*
[d] *Department of Mathematics, Faculty of Informatics and Control Systems, Georgian Technical University, 77, Kostava St., Tbilisi, Georgia*

Abstract

We derive criteria governing two-weight estimates for multilinear fractional integrals and appropriate maximal functions. The two and one weight problems for multi(sub)linear strong fractional maximal operators are also studied; in particular, we derive necessary and sufficient conditions guaranteeing the trace type inequality for this operator. We also establish the Fefferman-Stein type inequality, and obtain one-weight criteria when a weight function is of product type. As a consequence, appropriate results for multilinear Riesz potential operator with product kernels follow.

Keywords: Multilinear fractional integrals; multi(sub)linear strong fractional maximal operators; boundedness; two-weight inequality; trace inequality; Fefferman–Stein inequality.
2010 MSC: Primary 42B25, Secondary, 42B35

1. Introduction

Necessary and sufficient conditions governing the two-weight inequality for multilinear fractional integral operators are established. Our results involve, for example, criteria ensuring the trace type inequality and one-weight inequality for multi(sub)linear fractional maximal operators. Fefferman–Stein type inequality is also established for these operators. It should be also emphasized that appropriate results for multilinear Riesz potential operators follow.

Historically, multilinear fractional integrals were introduced in the papers by L. Grafakos [4], C. Kenig and E. Stein [7], L. Grafakos and N. Kalton [5]. In particular, these works deal with the operator

$$B_\alpha(f,g)(x) = \int\limits_{\mathbb{R}^n} \frac{f(x+t)g(x-t)}{|t|^{n-\alpha}}dt, \quad 0 < \alpha < n.$$

*This paper was publish in *Preprints of Abdus Salam School of Mathematical Sciences*, No. 481, at October 29, 2012 (see http://sms.edu.pk/journals/preprint/pre_481.pdf)

Email addresses: kokil@rmi.ge (Vakhtang Kokilashvili), mastylo@amu.edu.pl (Mieczyslaw Mastylo), alex72meskhi@yahoo.com; meskhi@rmi.ge (Alexander Meskhi)

In the mentioned papers it was proved that if $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$, where $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}$, then B_α is bounded from $L^{p_1} \times L^{p_2}$ to L^q.

As a tool to understand B_α, the operators

$$\mathcal{I}_\alpha(\overrightarrow{f})(x) = \int\limits_{(\mathbb{R}^n)^m} \frac{f_1(y_1)\cdots f_m(y_m)}{(|x-y_1|+\cdots+|x-y_m|)^{mn-\alpha}} d\overrightarrow{y},$$

where $x \in \mathbb{R}^n$, $0 < \alpha < nm$, $\overrightarrow{f} := (f_1, \cdots, f_m)$, $\overrightarrow{y} := (y_1, \cdots, y_m)$, were studied as well. Corresponding maximal operator is given by (see [15])

$$\mathcal{M}_\alpha(\overrightarrow{f})(x) = \sup_{Q \ni x} \prod_{i=1}^m \frac{1}{|Q|^{1-\alpha/(nm)}} \int\limits_Q |f_i(y_i)| dy_i, \quad 0 \le \alpha < mn,$$

where $|Q|$ denotes the volume of the cube Q with sides parallel to the coordinate axes.

This operator for $\alpha = 0$ was introduced and studied in [14].

It can be immediately checked that

$$\mathcal{I}_\alpha(\overrightarrow{f})(x) \ge c_n \mathcal{M}_\alpha(\overrightarrow{f})(x), \quad f_i \ge 0, \quad i = 1, \cdots, m.$$

In the sequel the following notation will be used:

$$\overrightarrow{p} := (p_1, \cdots, p_m), \quad \overrightarrow{w} = (w_1, \cdots, w_m),$$

where p_i are constants ($0 < p_i < \infty$) and w_i are a.e. positive functions defined on Euclidean space. It will be also assumed that

$$\frac{1}{p} = \sum_{i=1}^m \frac{1}{p_i}.$$

Throughout the paper we use the notation $\mathcal{Q}$ to denote the family of all cubes in $\mathbb{R}^n$ with sides parallel to the coordinate axes.

An almost everywhere positive and locally integrable function defined on $\mathbb{R}^n$ is called a weight.

Let $0 < r < \infty$ and let ρ be a weight on $\mathbb{R}^n$. We denote by $L^r_\rho(\mathbb{R}^n)$ a class of all measurable functions f on $\mathbb{R}^n$ such that

$$\|f\|_{L^r_\rho(\mathbb{R}^n)} := \Big(\int\limits_{\mathbb{R}^n} |f(x)|^r \rho(x) dx \Big)^{1/r} < \infty.$$

Definition 1.1 (vector Muckenhoupt condition)**.** Let $1 \le p_i < \infty$ for $i = 1, \cdots, m$. Let w_i be weights on $\mathbb{R}^n$, $i = 1, \cdots, m$. We say that $\overrightarrow{w} \in A_{\overrightarrow{p}}(\mathbb{R}^n)$ (or simply $\overrightarrow{w} \in A_{\overrightarrow{p}}$) if

$$\sup_{Q \in \mathcal{Q}} \Big(\frac{1}{|Q|} \int\limits_Q \prod_{i=1}^m w_i^{p/p_i} \Big)^{1/p} \prod_{i=1}^m \Big(\frac{1}{|Q|} \int\limits_Q w_i^{1-p_i'} \Big)^{1/p_i'} < \infty.$$

When $p_i = 1$, $\Big(\frac{1}{|Q|} \int\limits_Q w_i^{1-p_i'} \Big)^{1/p_i'}$ is understood as $(\inf_Q w_i)^{-1}$.

Remark 1.2. In the linear case ($m = 1$) the class $A_{\overrightarrow{p}}$ coincides with the well- known Muckenhoupt class A_p.

"Let us recall that, by definition, $A_\infty = \cup_{i=1}^{\infty} A_p$.

Further, let us recall that, by definition, $A_\infty = \cup_{i=1}^{\infty} A_p$.

Definition 1.3 (vector Muckenhoupt-Wheeden condition)**.** Let $1 \le p_i < \infty$ for $i = 1, \cdots, m$. Suppose that $p < q < \infty$. We say that $\overrightarrow{w} = (w_1, \cdots, w_m)$ satisfies $A_{\overrightarrow{p},q}(\mathbb{R}^n)$ condition ($\overrightarrow{w} \in A_{\overrightarrow{p},q}$) if

$$\sup_Q \Big(\frac{1}{|Q|}\int\limits_Q \Big(\prod_{i=1}^m w_i\Big)^q\Big)^{1/q} \prod_{i=1}^m \Big(\frac{1}{|Q|}\int\limits_Q w_i^{-p_i'}\Big)^{1/p_i'} < \infty.$$

Theorem A ([14]). *Let $1 < p_i < \infty$, $i = 1, \cdots, m$. Suppose that w_i are weights on $\mathbb{R}^n$. Then the operator $\mathcal{M}$ is bounded from $L^{p_1}_{w_1}(\mathbb{R}^n) \times \cdots \times L^{p_m}_{w_m}(\mathbb{R}^n)$ to $L^{p}_{\prod_{i=1}^m w_i^{p/p_i}}(\mathbb{R}^n)$ if and only if $\overrightarrow{w} \in A_{\overrightarrow{p}}(\mathbb{R}^n)$.*

Theorem B. ([15]) *Let $1 < p_1, \cdots, p_m < \infty$, $0 < \alpha < mn$, $\frac{1}{m} < p < \frac{n}{\alpha}$. Suppose that q is an exponent satisfying the condition $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$. Suppose that w_i are a.e. positive functions on $\mathbb{R}^n$ such that $w_i^{p_i}$ are weights. Then the inequality*

$$\Big(\int\limits_{\mathbb{R}^n} \Big(\big|\mathcal{N}_\alpha(\overrightarrow{f})(x)\big| \prod_{i=1}^m w_i(x)\Big)^q dx\Big)^{1/q} \le C \prod_{i=1}^m \Big(\int\limits_{\mathbb{R}^n} (|f_i(y_i)|w_i)^{p_i} dy_i\Big)^{1/p_i},$$

holds, where $\mathcal{N}_\alpha$ is $\mathcal{I}_\alpha$ or $\mathcal{M}_\alpha$, if and if $\overrightarrow{w} \in A_{\overrightarrow{p},q}(\mathbb{R}^n)$.

Regarding the two-weight result it is known the following result:

Theorem C. *Let $1 < p_1, \cdots, p_m < \infty$ and let $0 < \alpha < mn$. Let w_i and u be a.e. positive functions on $\mathbb{R}^n$. Further, assumed that q is an exponent satisfying the condition $\frac{1}{m} < p \le q < \infty$. Suppose also that*

(i) $q > 1$ and a pair $(u, \overrightarrow{w})$ satisfies the condition

$$\sup_Q |Q|^{\alpha/n+1/q-1/p}\Big(\frac{1}{|Q|}\int\limits_Q u^{qr}(x)dx\Big)^{1/qr} \prod_{i=1}^m \Big(\frac{1}{|Q|}\int\limits_Q w_i^{-p_i'r}(y_i)dy_i\Big)^{1/p_i'r} < \infty \tag{1}$$

for some $r > 1$;

(ii) $q \le 1$ and $(u, \overrightarrow{w})$ satisfies the condition

$$\sup_Q |Q|^{\alpha/n+1/q-1/p}\Big(\frac{1}{|Q|}\int\limits_Q u^{q}(x)dx\Big)^{1/qr} \prod_{i=1}^m \Big(\frac{1}{|Q|}\int\limits_Q w_i^{-p_i'r}(y_i)dy_i\Big)^{1/p_i'r} < \infty \tag{2}$$

for some $r > 1$. Then the inequality

$$\Big(\int\limits_{\mathbb{R}^n} (|\mathcal{I}_\alpha(\overrightarrow{f})(x)|u(x))^q dx\Big)^{1/q} \le c \prod_{i=1}^m \Big(\int\limits_{\mathbb{R}^n} (|f_i(x)|w_i(x))^{p_i} dx\Big)^{1/p_i}$$

holds for all $\overrightarrow{f} \in L^{p_1}_{w_1^{p_1}}(\mathbb{R}^n) \times \cdots \times L^{p_m}_{w_m^{p_m}}(\mathbb{R}^n)$.

Remark 1.4. The two-weight problem for sublinear fractional maximal operator

$$(M_\alpha f)(x) = \sup_{Q\ni x} \frac{1}{|Q|^{1-\alpha/n}} \int\limits_Q f(y)dy,$$

has been already solved. We mention the papers by E. Sawyer [19] for the conditions involving the operator itself and by R. L. Wheeden [24] for the Gabidzashvili-Kokilashvili type conditions (see also [9], [11] for the latter condition). For the solution of the two-weight problem for the Riesz fractional integral operator

$$(I_\alpha f)(x) = \int\limits_{\mathbb{R}^n} \frac{f(y)}{|x-y|^{n-\alpha}} dy$$

under different conditions we refer to the paper [20] and the monograph [11].

Remark 1.5. Conditions (1) and (2) are known as the "power bump" conditions (see [17]) for the linear case).

Remark 1.6. Some "power bump" sufficient conditions governing the two-weight inequality for $\mathcal{I}_\alpha$ were derived in [22].

Corollary A. *Let $0 < \alpha < mn$ and let $1 < p_i < \infty$, $i = 1, \cdots, m$. Assume that $\frac{1}{m} < p \le q < \infty$. Suppose that $u, w_1, \cdots w_m$ are a.e. positive functions on $\mathbb{R}^n$ such that $u^q, w_1^{-p_1'}, \cdots, w_m^{-p_m'} \in A_\infty(\mathbb{R}^n)$. Then the inequality*

$$\Big(\int\limits_{\mathbb{R}^n} \Big(|\mathcal{N}_\alpha(\overrightarrow{f})(x)|u(x)\Big)^q dx\Big)^{1/q} \le C \prod_{i=1}^m \Big(\int\limits_{\mathbb{R}^n} (|f_i(y_i)|w_i(y_i))^{p_i} dy_i\Big)^{1/p_i}, \tag{3}$$

where $\mathcal{N}_\alpha$ is $\mathcal{I}_\alpha$ or $\mathcal{M}_\alpha$ holds if and only if

$$\sup_{Q\in Q} |Q|^{\alpha/n+1/q-1/p} \Big(\frac{1}{|Q|}\int\limits_Q u^q(x)dx\Big)^{1/q} \prod_{i=1}^m \Big(\frac{1}{|Q|}\int\limits_Q w_i^{-p_i'}(y_i)dy_i\Big)^{1/p_i'} < \infty. \tag{4}$$

Theorem D. ([15]) *Let $0 \le \alpha < mn$ and let $1 \le p_i < \infty$, $1 = 1, \cdots, m$. Suppose that $\frac{1}{m} < p \le q < \infty$. Then the weak type inequality*

$$\|u\mathcal{M}_\alpha(\overrightarrow{f})\|_{L^{q,\infty}(\mathbb{R}^n)} \le c \prod_{i=1}^m \|w_i f_i\|_{L^{p_i}(\mathbb{R}^n)}$$

holds if and only if (4) *satisfied.*

The next statement shows that for $\mathcal{N}_\alpha = \mathcal{M}_\alpha$, only the "power bump" condition with respect to the right-hand side weight is sufficient for the strong–type inequality for $\mathcal{M}_\alpha$.

Theorem E. *Let $0 \le \alpha < mn$, $1 < p_1, \cdots, p_n$, $\frac{1}{m} < p \le q < \infty$. If $(u, \overrightarrow{w})$ satisfies* (2)*, then inequality* (3) *holds for $\mathcal{N}_\alpha = \mathcal{M}_\alpha$.*

Corollary B. *Let α, p_i, q satisfy the conditions of Theorem E. Suppose that $w_i^{-p_i'} \in A_\infty$. Then the inequality* (3) *holds for $\mathcal{N}_\alpha = \mathcal{M}_\alpha$ if and only if* (4) *holds.*

In [10], based on Corollary B, the authors of this paper characterized the inequality (3) for $\mathcal{N}_\alpha = \mathcal{I}_\alpha$ and $w_i \equiv$ *const* without any additional restriction on u under the D. Adams [1] type condition. In particular, they proved the next statement.

Theorem F. *Let $1 < p_i < \infty$, $i = 1, \cdots, m$. Assume that $\alpha < n/p$ and $p < q < \infty$. Then the following conditions are equivalent:*

(i) inequality (3) holds for $\mathcal{N}_\alpha = \mathcal{I}_\alpha$ and $w_i \equiv$ *const*;

(ii) the inequality

$$\|u\mathcal{M}_\alpha(\overrightarrow{f})\|_{L^{q,\infty}(\mathbb{R}^n)} \le c \prod_{i=1}^{m} \|f_i\|_{L^{p_i}(\mathbb{R}^n)}$$

holds;

(iii) the condition

$$\sup_{Q\in\mathcal{Q}} \Big(\int_Q u^q(x)dx \Big)^{1/q} |Q|^{\alpha-n/p} < \infty$$

is satisfied.

Remark 1.7. Finally we mention that Fefferman-Stein type inequality for the multi(sub)linear operator $\mathcal{M}_\alpha$ was studied in [18].

2. Preliminaries

Let $\mathcal{D}$ be the set of all dyadic cubes in $\mathbb{R}^n$. It is known that each $Q \in \mathcal{D}$ is the union of 2^n non-overlapping dyadic cubes (for the definition and some properties of the dyadic intervals see, for instance, [3], P. 136).

Definition 2.1. We say that a weight function ρ satisfies the dyadic reverse doubling condition ($\rho \in RD^{(d)}(\mathbb{R})$) if there exists a constant $d > 1$ such that

$$d\rho(Q') \le \rho(Q), \tag{5}$$

for all $Q', Q \in \mathcal{D}$, where $Q' \subset Q$ and $|Q| = 2^n|Q'|$.

Further, we say that a weight function ρ satisfies the reverse doubling condition ($\rho \in RD(\mathbb{R}^n)$) if (5) holds for all cubes Q', Q, where Q' is a subcube of Q arising dividing Q by 2^n equal parts.

We shall also need the next Carleson-Hörmander type embedding theorem regarding the dyadic intervals (see e.g., [21], [23]):

Theorem G. *Let $1 < r < q < \infty$ and let ρ be a weight function on $\mathbb{R}^n$ such that $\rho^{1-r'}$ satisfies the dyadic reverse doubling condition. Let $\{c_Q\}$ be non-negative numbers corresponding to dyadic cubes Q in $\mathbb{R}$. Then the following two statements are equivalent:*

(i) *There is a positive constant C such that*

$$\sum_{Q\in\mathcal{D}} c_Q \left(\frac{1}{|Q|} \int_Q g(x)dx \right)^q \le C \left(\int_{\mathbb{R}^n} g(x)^r \rho(x)dx \right)^{q/r}$$

for all non-negative $g \in L^r_\rho(\mathbb{R}^n)$;

(ii) *There is a positive constant* C_1 *such that*

$$c_Q \le C_1|Q|^q\left(\int_Q \rho(x)^{1-r'}dx\right)^{-q/r'}$$

for all $Q \in \mathcal{D}$.

This result yields the following corollary:

Corollary C. *Let* $1 < r < q < \infty$ *and let* ρ *be a weight function on* $\mathbb{R}^n$ *such that* $\rho^{1-r'}$ *satisfies the dyadic reverse doubling condition. Then the Carleson-Hörmander type inequality*

$$\sum_{Q\in\mathcal{D}}\left(\int_Q \rho^{1-r'}(x)dx\right)^{-q/r'}\left(\int_Q f(x)dx\right)^q \le c\left(\int_{\mathbb{R}^n} f^r(x)\rho(x)dx\right)^{q/r}$$

holds for all non-negative $f \in L^r_\rho(\mathbb{R}^n)$.

Let us recall some properties of vector Muckenhoupt condition:

Lemma 2.2 ([14], [15])**.** *Let* $1 < p_i < \infty$, $i = 1, \cdots, m$. *Then*

(i)

$$\prod_{i=1}^m A_{p_i}(\mathbb{R}^n) \subset A_{\overrightarrow{p}}(\mathbb{R}^n);$$

(ii) if $\overrightarrow{w} \in A_{\overrightarrow{p},q}(\mathbb{R}^n)$, where $1/q = 1/p - \alpha/n$, $0 < \alpha < mn$, then

$$\left(\prod_{i=1}^m w_i\right)^q \in A_{mq}(\mathbb{R}^n) \text{ and } w_i^{-p_i'} \in A_{mp_i'}(\mathbb{R}^n).$$

3. Main Results

Our aim in this paper is to improve the known results (see the statements above) regarding the two–weight strong-type inequality for $\mathcal{N}_\alpha$, where $\mathcal{N}_\alpha$ is $\mathcal{M}_\alpha$ or $\mathcal{I}_\alpha$, and to study the two-weight boundedness for the strong fractional maximal operator

$$\mathcal{M}^{(S)}_{\alpha_1,\cdots\alpha_k}(\overrightarrow{f})(x) = \sup_{Q^{(1)}\times\cdots\times Q^{(k)}\ni x} \frac{1}{|Q^{(1)}|^{m-\alpha_1/n}\times\cdots\times|Q^{(k)}|^{m-\alpha_k/n}}\prod_{i=1}^m \int\limits_{Q^{(1)}\times\cdots\times Q^{(k)}} |f_i(y_i)|dy_i, \qquad (6)$$

where $0 \le \alpha_1, \cdots, \alpha_k < mn$, $x = (x_1, \cdots, x_k) \in \mathbb{R}^{kn}$ $(k \ge 2)$ and the supremum is taken over all products of cubes $Q^{(1)}, \cdots, Q^{(k)} \in Q$ containing $x_1, \cdots, x_k$ respectively.

Operator (6) for $\alpha_1 = \cdots \alpha_k = 0$ and $n = 1$ was introduced in [6]. In this case we have multi(sub)linear strong maximal operator denoted by $\mathcal{M}^{(S)}$ and defined with respect to rectangles in $\mathbb{R}^k$ with sides parallel to the coordinate axes. In that paper the authors studied one and two–weight problems for $\mathcal{M}^{(S)}$. In particular, they proved that the one-weight inequality holds if and only if the weight satisfies the strong $A_{\overrightarrow{p}}$ condition ($A_{\overrightarrow{p}}$ condition written with respect to rectangles in $\mathbb{R}^k$).

Theorem 3.1. *Let $1 < p_i < \infty$, $i = 1, \cdots, m$. Suppose that $p < q < \infty$ and $0 < \alpha < mn$. Suppose also that $w_i^{-p_i'}$ satisfies the reverse doubling condition on $\mathbb{R}^n$, $i = 1, \cdots, m$. Then the inequality*

$$\|u\mathcal{M}_\alpha(\overrightarrow{f})\|_{L^q(\mathbb{R}^n)} \le C \prod_{i=1}^m \|w_i f_i\|_{L^{p_i}(\mathbb{R}^n)} \tag{7}$$

holds for all $\overrightarrow{f} \in L^{p_1}_{w_1^{p_1}}(\mathbb{R}^n) \times \cdots \times L^{p_m}_{w_m^{p_m}}(\mathbb{R}^n)$ if and only if condition (4) *is satisfied.*

Remark 3.2. Since the reverse doubling condition is weaker than the A_∞ condition (see e.g. [3]), Theorem 3.1 generalizes Corollary A for $\mathcal{N}_\alpha = \mathcal{M}_\alpha$.

Corollary 3.3. *Let conditions of Theorem 3.1 be satisfied and let, in addition, $u^q \in A_\infty(\mathbb{R}^n)$. Then the inequality*

$$\|u\mathcal{I}_\alpha(\overrightarrow{f})\|_{L^q(\mathbb{R}^n)} \le C \prod_{i=1}^m \|w_i f_i\|_{L^{p_i}(\mathbb{R}^n)}$$

holds for all $\overrightarrow{f} \in L^{p_1}_{w_1^{p_1}}(\mathbb{R}^n) \times \cdots \times L^{p_m}_{w_m^{p_m}}(\mathbb{R}^n)$ if and only if condition (4) *is satisfied.*

Corollary 3.4 ([15])**.** *Let $1 < p_i < \infty$ $(i = 1, \cdots, m)$ and let $p < q < \infty$. Suppose that $0 < \alpha < n$. We set $1/q = 1/p - \alpha/n$. Then the one-weight inequality*

$$\Big(\int_{\mathbb{R}^n} \Big[\Big|\mathcal{N}_\alpha(\overrightarrow{f})(x)\Big|\Big(\prod_{i=1}^m w_i(x)\Big)\Big]^q dx\Big)^{1/q} \le C \prod_{i=1}^m \Big(\int_{\mathbb{R}^n} \Big[\Big|f_i(x)\Big|w_i(x)\Big]^{p_i} dx\Big)^{1/p_i},$$

where $\mathcal{N}_\alpha$ is $\mathcal{M}_\alpha$ or $\mathcal{I}_\alpha$, holds if and only if $\overrightarrow{w} \in A_{\overrightarrow{p},q}$.

Regarding the strong maximal operator we have:

Theorem 3.5. *Let $1 < p_i < \infty$ $(i = 1, \cdots, m)$ and let $p < q < \infty$. Assume that $0 < \alpha_1, \cdots, \alpha_k < mn$. Suppose that u and w_i are a.e. positive functions defined on $\mathbb{R}^{kn}$ such that u^q, $w_1^{p_1}, \cdots, w_m^{p_m}$ are weights. Suppose, in addition, that w_i are of the product type: $w_i = w_i^{(1)} \times \cdots \times w_i^{(k)}$, $i = 1, \cdots, m$, where $w_i^{(1)}, \cdots, w_i^{(k)}$ are defined on $\mathbb{R}^n$. Assume also that $(w_i^{(1)})^{-p'}, \cdots, (w_i^{(k)})^{-p'} \in RD(\mathbb{R}^n)$. Then the inequality*

$$\|u\mathcal{M}^{(S)}_{\alpha_1,\cdots,\alpha_k}(\overrightarrow{f})\|_{L^q(\mathbb{R}^{kn})} \le C \prod_{i=1}^m \|w_i f_i\|_{L^{p_i}(\mathbb{R}^{kn})}$$

holds if and only if

$$\sup_{Q_1,\cdots,Q_k \in Q} |Q_1|^{\alpha_1/n-m} \times \cdots \times |Q_k|^{\alpha_k/n-m} \Big(\int_{Q_1\times\cdots\times Q_k} u^q(x)dx\Big)^{1/q} \prod_{i=1}^m \Big(\int_{Q_1\times\cdots\times Q_k} w_i^{-p_i'}(y_i)dy_i\Big)^{1/p_i'} < \infty. \tag{8}$$

Since for a constant weight function the reverse doubling condition is automatically satisfied, from Theorem 3.5 we have a characterization of the boundedness for the operator $\mathcal{M}_{\alpha_1,\cdots,\alpha_k}$ from L^p to $L^q_{u^q}$. In particular, we have

Corollary 3.6. *Let $p, q, \alpha_1, \cdots, \alpha_k$ satisfy the conditions of Theorem 3.5. Then the inequality*

$$\|u\mathcal{M}^{(S)}_{\alpha_1,\cdots,\alpha_k}(\overrightarrow{f})\|_{L^q(\mathbb{R}^{kn})} \le c\prod_{i=1}^{m}\|f_i\|_{L^{p_i}(\mathbb{R}^{kn})}$$

holds for all $\overrightarrow{f} \in L^{p_1}(\mathbb{R}^{kn}) \times \cdots \times L^{p_m}(\mathbb{R}^{kn})$ if and only if

$$\sup_{Q_1,\cdots,Q_k}\Big(\int\limits_{Q_1\times\cdots\times Q_k} u^q(x)dx\Big)|Q_1|^{q(\alpha_1/n-1/p)}\times\cdots\times|Q_k|^{q(\alpha_k/n-1/p)} < \infty, \tag{9}$$

where, as before, $\frac{1}{p} = \sum_{i=1}^{m}\frac{1}{p_i}$.

Similar results were derived in [12] (see also [13], Ch.4) for sublinear maximal operators.

It is natural to study the appropriate potential operator with product kernels:

$$\mathcal{I}_{\alpha_1,\cdots,\alpha_k}(\overrightarrow{f})(x^{(1)},\cdots,x^{(k)}) = \int\limits_{(\mathbb{R}^{kn})^m}\frac{\prod_{i=1}^{m} f_i(y_i^{(1)},\cdots,y_i^{(k)})dy_1^{(1)}\cdots dy_k^{(1)}\cdots dy_1^{(m)},\cdots dy_k^{(m)}}{\prod_{s=1}^{k}(\sum_{i=1}^{m}|x^{(s)}-y_i^{(s)}|)^{mn-\alpha_s}},$$

where $x^{(i)} \in \mathbb{R}^n$, $i = 1, \cdots, m$.

It is easy to see that the following pointwise inequality holds

$$\mathcal{I}_{\alpha_1,\cdots,\alpha_k}(\overrightarrow{f}) \ge c_n\mathcal{M}^{(S)}_{\alpha_1,\cdots,\alpha_k}(\overrightarrow{f}), \quad f_i \ge 0,\ i = 1,\cdots,m.$$

If $m = 1$, the operator $\mathcal{I}_{\alpha_1,\cdots,\alpha_n}$ is the linear Riesz potential operator with product kernels denoted by $I_{\alpha_1,\cdots,\alpha_n}$. The one-weight criteria for this operator were established in [8], while the trace type inequality was derived in [12] (see also [13], Ch.4).

Theorem 3.7. *Let $p_i, q, \alpha_1, \cdots, \alpha_k$ satisfy the conditions of Theorem 3.5. Suppose that u^q satisfies $A_\infty(\mathbb{R}^n)$ condition in each variable separately uniformly to another one. Then the trace type inequality*

$$\|u\mathcal{I}_{\alpha_1,\cdots\alpha_k}(\overrightarrow{f})\|_{L^q(\mathbb{R}^{kn})} \le c\prod_{i=1}^{m}\|w_i f_i\|_{L^{p_i}(\mathbb{R}^{kn})}$$

holds if and only if (8) *is true.*

Corollary 3.8 (One-weight inequality)**.** *Let $1 < p_i < \infty$, $i = 1, \cdots, m$. Suppose that $p < q < \infty$ and $0 < \alpha < mn$. We set $1/q = 1/p - \alpha/n$. Suppose that w_i are a.e. positive functions such that $w_i^{p_i}$ are weights on $\mathbb{R}^{kn}$ and $w_i = w_i^{(1)} \times \cdots \times w_i^{(k)}$, $i = 1, \cdots, m$, where $w_i^{(j)}$ are defined on $\mathbb{R}^n$. Then the one-weight inequality*

$$\Big(\int\limits_{\mathbb{R}^{nk}}\Big[\Big|\mathcal{S}_\alpha(\overrightarrow{f})(x)\Big|\Big(\prod_{i=1}^{m}w_i(x)\Big)\Big]^q dx\Big)^{1/q} \le C\prod_{i=1}^{m}\Big(\int\limits_{\mathbb{R}^{nk}}\Big[\Big|f_i(x)\Big|w_i(x)\Big]^{p_i}dx\Big)^{1/p_i},$$

where $\mathcal{S}_\alpha$ is $\mathcal{M}^{(S)}_{\alpha,\cdots,\alpha}$ or $\mathcal{I}_{\alpha,\cdots,\alpha}$, holds if and only if $\overrightarrow{w} \in \mathcal{A}_{\overrightarrow{P},q}(\mathbb{R}^{kn})$, i.e.

$$\sup_{Q_1,\cdots,Q_k\in\mathcal{Q}}\Big(\frac{1}{\prod_{i=1}^{k}|Q_i|}\int\limits_{Q_1\times\cdots\times Q_k}\Big(\prod_{i=1}^{m}w_i\Big)^q\Big)^{1/q}\prod_{i=1}^{m}\Big(\frac{1}{\prod_{i=1}^{k}|Q_i|}\int\limits_{Q_1\times\cdots\times Q_k}w_i^{-p_i'}\Big)^{1/p_i'} < \infty, \tag{10}$$

where the supremum is taken over all cubes $Q_j \subset \mathbb{R}^n$, $j = 1, \cdots, k$.

Our next result concerns with the Fefferman-Stein type inequality for the (sub)linear strong fractional maximal operator $\mathcal{M}^{(S)}_{\alpha_1,\cdots,\alpha_k}$.

Theorem 3.9. *Let $p, q, \alpha_1, \cdots, \alpha_k$ satisfy the conditions of Theorem 3.5. Suppose that v is an a.e. positive function on $\mathbb{R}^{kn}$. Then there is a positive constant C such that*

$$\|\mathcal{M}^{(S)}_{\alpha_1,\cdots,\alpha_k}(\overrightarrow{f})\|_{L^q_v(\mathbb{R}^{kn})} \leq c \prod_{i=1}^{m} \|(\overline{M}_{\alpha_1,\cdots,\alpha_k} v)^{p/(p_i q m)} f_i\|_{L^{p_i}(\mathbb{R}^{kn})}$$

holds, where $\overline{M}_{\alpha_1,\cdots,\alpha_m}$ is the sublinear strong fractional maximal operator defined on $\mathbb{R}^{kn}$ and given by

$$\left(\overline{M}_{\alpha_1,\cdots,\alpha_m} v\right)(x^{(1)},\cdots,x^{(k)})$$

$$= \sup_{Q_1 \ni x^{(1)},\cdots,Q_k \ni x^{(k)}} |Q_1|^{q\left(\frac{\alpha_1}{n}-\frac{1}{p}\right)} \times \cdots \times |Q_k|^{q\left(\frac{\alpha_k}{n}-\frac{1}{p}\right)} \int_{Q_1 \times,\cdots,\times Q_k} v(y^{(1)},\cdots,y^{(k)}) dy^{(1)} \cdots dy^{(k)}.$$

4. Proofs of the Main Results

In this section we give proofs of the main results of this paper. First we formulate the following auxiliary statement:

Lemma 4.1. *[15] Let $\mathcal{M}^k_\alpha$ be defined as $\mathcal{M}_\alpha$ but over cubes with side length less than or equal to 2^k. Suppose that $Q_k := [-2^{k+2}, 2^{k+2})^n$, $\tau_t g(x) := g(x-t)$ and $\overrightarrow{\tau}_t(\overrightarrow{f}) := (\tau_t f_1, \cdots, \tau_t f_m)$. Then there is a positive constant c depending only on n, m, α and q such that for each $k \in \mathbb{Z}$, $\overrightarrow{f}$ and every $x \in \mathbb{R}^n$,*

$$(\mathcal{M}^k_\alpha(\overrightarrow{f})(x))^q \leq \frac{c}{|Q_k|} \int_{Q_k} \left[\left(\tau_{-t} \circ \mathcal{M}^{(d)}_\alpha \circ \overrightarrow{\tau}_t\right)(\overrightarrow{f})(x)\right]^q dt \tag{11}$$

Proof of Theorem 3.1 . Necessity. Let us first show that

$$\int_Q w_i^{-p_i'}(y_i) dy_i < \infty, \tag{12}$$

for all $i = 1, \cdots, m$ and all cubes Q. Indeed, suppose that

$$\int_Q w_{i_1}^{-p_{i_1}'}(y_{i_1}) dy_{i_1} = \cdots = \int_Q w_{i_k}^{-p_{i_k}'}(y_{i_k}) dy_{i_k} = \infty, \tag{13}$$

for some $1 \leq i_1 < \cdots < i_k \leq m$. Then by the duality arguments, we have that there are non–negative functions $g_{i_1}, \cdots g_{i_k}$ such that

$$\int_Q w_{i_j}^{-1} g_{i_j} = \infty, \quad j = 1, \cdots, k,$$

and $g \in L^{p_{i_j}}(Q)$, $j = 1, \cdots, k$. Let us assume that $\overrightarrow{f} = (f_1, \cdots, f_m)$, where $f_j = w_{i_j}^{-1} g_{i_j} \chi_Q$, $j = 1, \cdots, k$ and $f_n = \chi_Q$ for other n. Then

$$\|u\mathcal{M}_\alpha(\overrightarrow{f})\|_{L^q(\mathbb{R}^n)} \geq \frac{1}{|Q|^{m-\alpha/n}}\Big(\prod_{j=1}^{k}\int\limits_Q w_{i_j}^{-1}g_{i_j}\Big)|Q|^{m-1}\|u\|_{L^q(Q)} = \infty.$$

On the other hand,

$$\prod_{i=1}^{m}\|f_i w_i\|_{L^{p_i}(\mathbb{R})} < \infty$$

because $w_i^{p_i}$ are locally integrals and $g_{i_j} \in L^{p_{i_j}}(\mathbb{R}^n)$, $j = 1, \cdots, k$.

Sufficiency. First we show that the two-weight inequality

$$\|u\mathcal{M}_\alpha^{(d)}(\overrightarrow{f})\|_{L^q(\mathbb{R}^n)} \leq C\prod_{i=1}^{m}\|w_i f_i\|_{L^{p_i}(\mathbb{R}^n)} \tag{14}$$

holds if (4) holds provided that $w_i^{-p_i'} \in RD^{(d)}(\mathbb{R}^n)$, $i = 1, \cdots, m$, where $\mathcal{M}_\alpha^{(d)}$ is a dyadic multi(sub)linear fractional maximal operator defined by

$$(\mathcal{M}_\alpha^{(d)}\overrightarrow{f})(x) = \sup_{Q\ni Q, Q\ni x}\prod_{i=1}^{m}\frac{1}{|Q|^{1-\alpha/(nm)}}\int\limits_Q |f_i(y_i)|dy_i, \quad 0 < \alpha < mn.$$

To show that (4) implies (14) we argue as follows: for every $x \in \mathbb{R}^n$ let us take $Q_x \in \mathcal{D}$ such that $Q_x \ni x$ and

$$(\mathcal{M}_\alpha^{(d)}\overrightarrow{f})(x) \leq \frac{2}{|Q_x|^{m-\alpha/n}}\prod_{i=1}^{m}\int\limits_{Q_x}|f_i(y_i)|dy_i. \tag{15}$$

Without loss of generality we can assume, for example, that f_i, $i = 1, \cdots, m$ are non–negative, bounded and have compact supports.

Let us introduce a set

$$F_Q = \{x \in \mathbb{R}^n : x \in Q \text{ and } (15) \ \textit{holds for} \ Q\}.$$

It is obvious that $F_Q \subset Q$ and $\mathbb{R}^n = \cup_{Q\in\mathcal{D}}F_Q$.

Further, we have

$$\int\limits_{\mathbb{R}^n}(\mathcal{M}_\alpha^{(d)}\overrightarrow{f})(x))^q(x)u^q(x)dx \leq \sum_{Q\in\mathcal{D}}\int\limits_{F_Q}(\mathcal{M}_\alpha^{(d)}\overrightarrow{f})^q u^q(x)dx$$

$$\leq 2^q\sum_{Q\in\mathcal{D}}|Q|^{(\alpha/n-m)q}\Big(\int\limits_Q u^q(x)dx\Big)\Big(\prod_{i=1}^{m}\int\limits_Q f_i(y_i)dy_i\Big)^q$$

$$\leq c\sum_{Q\in\mathcal{D}}\Big(\prod_{i=1}^{m}\int\limits_Q w_i^{-p'}(y_i)dy_i\Big)^{-q/p_i'}\Big(\prod_{i=1}^{m}\int\limits_Q f_i(y_i)dy_i\Big)^q$$

(using Hölder's inequality $\sum_k a_k^{(1)} \times \cdots \times a_k^{(m)} \leq \prod_{j=1}^m(\sum_k(a_k^{(j)})^{p_j/p})^{p/p_j}$ for positive sequences $\{a_k^{(j)}\}$, $j = 1, \cdots, m$)

$$\le \Big[\sum_{Q\in\mathcal{D}} \Big(\int_Q w_1^{-p_1'}(y_1)dy_1 \Big)^{-(qp_1)/(pp_1')} \Big(\int_Q f_1(y_1)dy_1 \Big)^{qp_1/p} \Big)^{p/p_1}$$

$$\times \cdots \times \Big(\sum_{Q\in\mathcal{D}} \Big(\int_Q w_m^{-p_m'}(y_m)dy_m \Big)^{-(qp_m)/(pp_m')} \Big(\int_Q f_m(y_m)dy_m \Big)^{qp_1/p} \Big]^{p/p_1}$$

(Applying Theorem G for the exponents $(p_i, qp_i/p)$, $i = 1, \cdots, m$)

$$\le c\| f_1 w_1\|^q_{L^{p_1}(\mathbb{R}^n)} \times \cdots \times \| f_m w_m\|^q_{L^{p_m}(\mathbb{R}^n)}.$$

Now we pass from $\mathcal{M}_\alpha^{(d)}$ to $\mathcal{M}_\alpha$. By using (11) we see that

$$\|u\mathcal{M}_\alpha(\overrightarrow{f})\|_{L^q(\mathbb{R}^n)} \le \sup_t \|u\Big(\tau_{-t} \circ \mathcal{M}_\alpha^{(d)} \circ \overrightarrow{\tau}_t\Big)(\overrightarrow{f})\|_{L^q(\mathbb{R}^n)}.$$

Observe now that a pair $(\tau_t u, \overrightarrow{\tau}_t \overrightarrow{w})$ satisfy condition (4) independent of t. Notice also that the weight functions $\tau_t \circ w_i$ satisfy the reverse doubling condition. Hence,

$$\|u\Big(\tau_{-t} \circ \mathcal{M}_\alpha^{(d)} \circ \overrightarrow{\tau}_t\Big)(\overrightarrow{f})\|_{L^q(\mathbb{R}^n)} \le c \prod_{i=1}^m \|w_i f_i\|_{L^{p_i}(\mathbb{R}^n)}$$

with the constant c independent of t. Finally, we conclude that

$$\int_{\mathbb{R}^n} (\mathcal{M}_\alpha \overrightarrow{f})(x))^q(x)u^q(x)dx \le \sup_t \Big\|u\Big(\tau_{-t} \circ \mathcal{M}_\alpha^{(d)} \circ \overrightarrow{\tau}_t\Big)(\overrightarrow{f})\Big\|_{L^q(\mathbb{R}^n)} \le c \prod_{i=1}^m \| f_i w_i\|_{L^{p_i}(\mathbb{R}^n)}.$$

□

Lemma 4.2. *Let $0 < q < \infty$ and let $0 < \alpha < mn$. Suppose that a weight function v satisfies $A_\infty(\mathbb{R}^n)$. Then there is a positive constant C such that for all $\overrightarrow{f}$ the inequality*

$$\int_{\mathbb{R}^n} |\mathcal{I}_\alpha(\overrightarrow{f})(x)|^q v(x)dx \le c \int_{\mathbb{R}^n} (\mathcal{M}_\alpha(\overrightarrow{f})(x))^q v(x)dx$$

holds.

Corollary 3.3 follows immediately from Theorem 3.1 and Lemma 4.2.

Corollary 3.4 for $\mathcal{M}_\alpha$ is a direct consequence of Lemma 2.2 and the fact that the Muckenhoupt condition implies the reverse doubling condition (see e.g. [3], [24]). Now the result for the operator $\mathcal{I}_\alpha$ follows by applying Lemma 4.2

To prove Theorem 3.4 we introduce a strong dyadic fractional operator

$$(\mathcal{M}_{\alpha,\beta}^{(S),(d)}(\overrightarrow{f})(x) = \sup_{Q^{(1)}\times Q^{(2)}\ni x,\ Q^{(1)},Q^{(2)}\in\mathcal{D}} \frac{1}{|Q^{(1)}|^{m-\alpha/n}|Q^{(2)}|^{m-\beta/n}} \prod_{i=1}^m \int_{Q^{(1)}\times Q^{(2} |f_i(y_i)|dy_i.$$

We need also the following notation

$$(\mathcal{M}^{(S),(k)}_{\alpha,\beta}(\overrightarrow{f})(x) = \sup_{Q^{(1)}\times Q^{(2)}\ni x,|Q^{(1)}|,|Q^{(2)}|\le 2^k} \frac{1}{|Q^{(1)}|^{m-\alpha/n}|Q^{(2)}|^{m-\beta/n}} \prod_{i=1}^{m}\int_R |f_i(y_i)|dy_i; \tag{16}$$

$$Q_k := [-2^{k+2}, 2^{k+2})^{2n}, \quad \tau_{t,\delta}g(x^{(1)}, x^{(2)}) := g(x^{(1)} - t, x^{(2)} - \delta)$$

$$\overrightarrow{\tau}_{t,\delta}(\overrightarrow{f}) := (\tau_{t,\delta}f_1, \cdots, \tau_{t,\delta}f_m),$$

where g and f are measurable functions on $\mathbb{R}^{2n}$.

Lemma 4.3. *There exists a positive constant c depending only on n, m, α, β and q such that for all $k \in \mathbb{Z}$, $\overrightarrow{f}$ and every $x \in \mathbb{R}^{2n}$,*

$$(\mathcal{M}^{(S),(k)}_{\alpha,\beta}(\overrightarrow{f})(x))^q \le \frac{c}{|Q_k|^2}\int_{Q_k^2}\Big[\Big(\tau_{-t,-\delta}\circ\mathcal{M}^{(S),(d)}_{\alpha,\beta}\circ\overrightarrow{\tau}_{t,\delta}\Big)(\overrightarrow{f})(x)\Big]^q dtd\delta. \tag{17}$$

Proof of this lemma follows in the same way as in the case of (sub)linear case (see [12], [13], P.); therefore we omit the details.

Proof of Theorem 3.5. For simplicity we prove the theorem for $k = 2$. The proof for other k is similar.

Necessity follows in the same way as in the case of the operator $\mathcal{M}_\alpha$.

To show *Sufficiency* we assume that f_i are non-negative and bounded with compact supports. For every $(x^{(1)}, x^{(2)}) \in \mathbb{R}^{2n}$, we take $Q^{(1)}, Q^{(2)} \in \mathcal{D}$ such that $Q^{(1)} \ni x^{(1)}$, $Q^{(2)} \ni x^{(2)}$ and

$$\mathcal{M}^{(S),(d)}_{\alpha_1,\alpha_2}(\overrightarrow{f})(x^{(1)}, x^{(2)}) \le \frac{2}{|Q^{(1)}|^{m-\alpha_1/n}|Q^{(1)}|^{m-\alpha_2/n}}\prod_{i=1}^{m}\int_{Q^{(1)}\times Q^{(2)}} f_i(y_i^{(1)}, y_i^{(2)})dy_i^{(1)}dy_i^{(2)}. \tag{18}$$

We construct the set:

$$F_{Q^{(1)},Q^{(2)}} := \{x = (x^{(1)}, x^{(2)}) \in \mathbb{R}^{2n} : x^{(1)} \in Q^{(1)}, x^{(2)} \in Q^{(2)} \text{and (18) holds for } Q^{(1)} \text{ and } Q^{(2)}\}.$$

Since $F_{Q^{(1)},Q^{(2)}} \subset Q^{(1)} \times Q^{(2)}$, arguing as in the proof of Theorem 3.1 we find that

$$\int_{\mathbb{R}^{2n}} (\mathcal{M}^{(d)}_{\alpha_1,\alpha_2}(\overrightarrow{f}))(x^{(1)}, x^{(2)}))^q u^q(x^{(1)}, x^{(2)})dx^{(1)}x^{(2)}$$

$$\le \sum_{Q_1,Q_2\in\mathcal{D}}\int_{F_{Q^{(1)},Q^{(2)}}} (\mathcal{M}^{(d)}_{\alpha_1,\alpha_2}(\overrightarrow{f})(x^{(1)}, x^{(2)}))^q u^q(x^{(1)}, x^{(2)})dx^{(1)}x^{(2)}$$

$$\le 2^q \sum_{Q^{(1)},Q^{(2)}\in\mathcal{D}} |Q|^{(\alpha/n-m)q}\Big(\int_Q u^q(x^{(1)}, x^{(2)})dx^{(1)}x^{(2)}\Big)\Big(\prod_{i=1}^{m}\int_{Q^{(1)}\times Q^{(2)}} f_i(y_i^{(1)}, y_i^{(2)})dy_i^{(1)}dy_i^{(2)}\Big)^q$$

$$\le c \sum_{Q^{(1)},Q^{(2)}\in\mathcal{D}} \prod_{i=1}^{m} \Big(\int\limits_{Q^{(1)}\times Q^{(2)}} w_i^{-p'}(y_i)dy_i \Big)^{-q/p_i'} \Big(\prod_{i=1}^{m} \int\limits_{Q} f_i(y_i^{(1)}, y_i^{(2)})dy_i^{(1)}dy_i^{(2)} \Big)^{q}$$

(due to Hölder's inequality $\sum_k a_k^{(1)}\times\cdots\times a_k^{(m)} \le \prod_{j=1}^{m} (\sum_k (a_k^{(j)})^{p_j/p})^{p/p_j}$ for positive sequences $\{a_k^{(j)}\}$, $j = 1,\cdots,m$)

$$\le \Big[\sum_{Q^{(1)},Q^{(2)}\in\mathcal{D}} \Big(\int\limits_{Q^{(1)}} (w_1^{(1)})^{-p_1'}(y_1^{(1)})dy_1^{(1)} \Big)^{-(qp_1)/(pp_1')} \Big(\int\limits_{Q^{(2)}} (w_1^{(2)})^{-p_1'}(y_1^{(2)})dy_1^{(2)} \Big)^{-(qp_1)/(pp_1')}$$

$$\times \Big(\int\limits_{Q^{(1)}} \Big(\int\limits_{Q^{(2)}} f_1(y_1^{(1)}, y_1^{(2)})dy_1^{(1)} \Big) dy_1^{(2)} \Big)^{qp/p_1} \Big]^{p_1/p}$$

$$\times\cdots\times \Big[\sum_{Q^{(1)},Q^{(2)}\in\mathcal{D}} \Big(\int\limits_{Q^{(1)}} (w_m^{(1)})^{-p_m'}(y_m^{(1)})dy_m^{(1)} \Big)^{-(qp_m)/(pp_m')} \Big(\int\limits_{Q^{(2)}} (w_m^{(2)})^{-p_1'}(y_m^{(2)})dy_m^{(2)} \Big)^{-(qp_1)/(pp_m')}$$

$$\times \Big(\int\limits_{Q^{(1)}} \Big(\int\limits_{Q^{(2)}} f_1(y_m^{(1)}, y_m^{(2)})dy_m^{(1)} \Big)\, dy_m^{(2)} \Big)^{qp/p_m} \Big]^{p_m/p}$$

(by virtue of Theorem G for the exponents $(p_i, qp_i/p)$, $i = 1,\cdots,m$, with respect to the first variable)

$$\le \Big[\sum_{Q^{(2)}\in\mathcal{D}} \Big(\int\limits_{Q^{(2)}} (w_1^{(2)})^{-p_1'}(y_1^{(2)})dy_1^{(2)} \Big)^{-(qp_1)/(pp_1')} \Big(\int\limits_{\mathbb{R}^n} (w_1^{(1)})^{p_1}(y_1^{(1)}) \Big(\int\limits_{Q^{(2)}} f_1(y_1^{(1)}, y_1^{(2)})dy_1^{(2)} \Big)^{p_1} dy_1^{(1)} \Big)^{q/p} \Big]^{p/p_1}$$

$$\times\cdots\times \Big[\sum_{Q^{(2)}\in\mathcal{D}} \Big(\int\limits_{Q^{(2)}} (w_m^{(2)})^{-p_m'}(y_m^{(2)})dy_m^{(2)} \Big)^{-(qp_m/pp_m')} \Big(\int\limits_{\mathbb{R}^n} (w_m^{(1)})^{p_m}(y_m^{(1)}) \Big(\int\limits_{Q^{(2)}} f_m(y_m^{(1)}, y_m^{(2)})dy_1^{(2)} \Big)^{p_m} dy_m^{(1)} \Big)^{q/p} \Big]^{p/p_m}$$

(using generalized Minkowski's inequality)

$$\le \Big[\sum_{Q^{(2)}\in\mathcal{D}} \Big(\int\limits_{Q^{(2)}} (w_1^{(2)})^{-p_1'}(y_1^{(2)})dy_1^{(2)} \Big)^{-(qp_1/pp_1')} \Big(\int\limits_{Q^{(2)}} \Big(\int\limits_{\mathbb{R}^n} f_1^{p_1}(y_1^{(1)}, y_1^{(2)})(w_1^{(1)})^{p_1}(y_1^{(1)})dy_1^{(1)} \Big)^{1/p_1} dy_1^{(2)} \Big)^{qp_1/p} \Big]^{p/p_1}$$

$$\times\cdots\times \Big[\sum_{Q^{(2)}\in\mathcal{D}} \Big(\int\limits_{Q^{(2)}} (w_m^{(2)})^{-p_m'}(y_m^{(2)})dy_m^{(2)} \Big)^{-(qp_m/pp_m')}$$

$$\times \Big(\int\limits_{Q^{(2)}} \Big(\int\limits_{\mathbb{R}^n} f_m^{p_m}(y_m^{(1)}, y_m^{(2)})(w_m^{(1)})^{p_m}(y_m^{(1)})dy_m^{(1)} \Big)^{1/p_m} dy_m^{(2)} \Big)^{qp_m/p} \Big]^{p/p_m}$$

(by virtue of Theorem G for the pairs of exponents: $(p_i, qp_i/p)$, $i = 1, \cdots, m$ with respect to the second variable)

$$\leq c\|f_1 w_1\|^q_{L^{p_1}(\mathbb{R}^{2n})} \times \cdots \times \|f_m w_m\|^q_{L^{p_m}(\mathbb{R}^{2n})}.$$

Arguing now as in the proof of Theorem 3.1 and using Lemma 4.3 we can pass from the dyadic strong maximal function $\mathcal{M}^{(S)(d)}_{\alpha_1,\alpha_2}$ to the operator $\mathcal{M}^{(S)}_{\alpha_1,\alpha_2}$ and get the desired result.

□

Theorem 3.7 follows from Theorem 3.5 and by using Lemma 4.2 with respect to each variable separately.

Proof of Corollary 3.8. We assume that $k = 2$ because the proof for higher k is similar. Suppose that $w_i = w_i^{(1)} w_i^{(2)}$, where $w_i^{(1)}$ and $w_i^{(2)}$ are defined on $\mathbb{R}^n$. Assume that $\mathcal{S}_{\alpha_1,\alpha_2} = \mathcal{M}_{\alpha_1,\alpha_2}$. First observe that condition (10) implies that

$$\sup_{Q_1 \in Q} \Big(\frac{1}{|Q_1|}\int_{Q_1} \Big[\prod_{i=1}^m w_i^{(1)}(x^{(1)})\Big]^q dx^{(1)}\Big)^{1/q} \prod_{i=1}^m \Big(\frac{1}{|Q_1|}\int_{Q_1} (w_i^{(1)})^{-p_i'}(x^{(1)})dx^{(1)}\Big)^{1/p'} < \infty,$$

and

$$\sup_{Q_2 \in Q} \Big(\frac{1}{|Q_2|}\int_{Q_2} \Big[\prod_{i=1}^m w_i^{(2)}(x^{(2)})\Big]^q dx^{(2)}\Big)^{1/q} \prod_{i=1}^m \Big(\frac{1}{|Q_2|}\int_{Q_2} (w_i^{(2)})^{-p_i'}(x^{(2)})dx^{(2)}\Big)^{1/p_i'} < \infty.$$

Consequently, by Lemma 2.2 we have that $(w_i^{(1)})^{-p_i'}, (w_i^{(2)})^{-p_i'} \in A_{mp_i'}(\mathbb{R}^n)$. Hence, (see e.g., [3] or [24]) $(w_i^{(1)})^{-p_i'}, (w_i^{(2)})^{-p_i'}$ satisfy the reverse doubling condition on $\mathbb{R}^n$. Now the result follows from Theorem 3.5.

The proof for $\mathcal{S}_{\alpha_1,\alpha_2} = \mathcal{I}_{\alpha_1,\alpha_2}$, follows from Lemmas 2.2 and 4.2.

□

Proof of Theorem 3.9. As before it is enough to prove the inequality

$$\|\mathcal{M}^{(S),(d)}_{\alpha,\beta}(\overrightarrow{f})\|_{L^q_v(\mathbb{R}^{2n})} \leq c\prod_{i=1}^m \|(\overline{M}_{\alpha,\beta} v)^{p/(p_i q m)} f_i\|_{L^{p_i}(\mathbb{R}^{2n})},$$

where $\mathcal{M}^{(S),(d)}_{\alpha_1,\alpha_2}$ is a dyadic strong fractional (sub)linear maximal operator defined above. Constructing the sets $F_{Q^{(1)},Q^{(2)}}$ for dyadic cubes $Q^{(1)}$ and $Q^{(2)}$ (see the proof of Theorem 3.5) we have

$$\int_{\mathbb{R}^{2n}} \Big(\mathcal{M}^{(S),(d)}_{\alpha_1,\alpha_2}(\overrightarrow{f})(x^{(1)}, x^{(2)})\Big)^q v(x^{(1)}, x^{(2)})dx^{(1)}dx^{(2)} \leq 2^q \sum_{Q^{(1)},Q^{(2)} \in \mathcal{D}} \Big(\int_{Q^{(1)}\times Q^{(2)}} v(x^{(1)}, x^{(2)})dx^{(1)}dx^{(2)}\Big)$$

$$\times |Q^{(1)}|^{(\alpha_1/n-m)q}|Q^{(2)}|^{(\alpha_2/n-m)q} \prod_{i=1}^m \Big(\int_{Q^{(1)}\times Q^{(2)}} |f_i(y_i^{(1)}, y_i^{(2)})|dy_i^{(1)}dy_i^{(2)})\Big)^q$$

$$= 2^q \sum_{Q^{(1)},Q^{(2)} \in \mathcal{D}} |Q^{(1)}|^{(\alpha_1/n-m)q}|Q^{(2)}|^{(\alpha_2/n-m)q}$$

$$\times\prod_{i=1}^{m}\Big(\int\limits_{Q^{(1)}\times Q^{(2)}}|f_i(y_i^{(1)},y_i^{(2)})|\Big(\int_{Q^{(1)}\times Q^{(2)}}v(x^{(1)},x^{(2)})dx^{(1)}dx^{(2)}\Big)^{1/(qm)}dy_i^{(1)}dy_i^{(2)})\Big)^q$$

$$\le c\sum_{Q^{(1)},Q^{(2)}\in\mathcal{D}}|Q^{(1)}|^{q(1/p-m)}|Q^{(2)}|^{(1/p-m)}\prod_{i=1}^{m}\Big(\int\limits_{Q^{(1)}\times Q^{(2)}}|f_i(y_i^{(1)},y_i^{(2)})|(\overline{M}_{\alpha_1,\alpha_2}(v)(y_i^{(1)},y_i^{(2)}))^{1/(qm)}dy_i^{(1)}dy_i^{(2)})\Big)^q$$

(using the fact that $|Q^{(1)}\times Q^{(2)}|^{q(1/p-m)}=\prod_{i=1}^{m}|Q^{(1)}\times Q^{(2)}|^{-q/p_i'}$)

$$=c\sum_{Q^{(1)},Q^{(2)}\in\mathcal{D}}\prod_{i=1}^{m}|Q^{(1)}|^{-q/p_i'}|Q^{(2)}|^{-q/p_i'}$$

$$\times\prod_{i=1}^{m}\Big(\int\limits_{Q^{(1)}\times Q^{(2)}}|f_i(y_i^{(1)},y_i^{(2)})|(\overline{M}_{\alpha_1,\alpha_2}(v)(y_i^{(1)},y_i^{(2)}))^{1/(qm)}dy_i^{(1)}dy_i^{(2)})\Big)^q$$

(by virtue of Hölder's inequality)

$$\le c\Big[\sum_{Q^{(1)},Q^{(2)}\in\mathcal{D}}|Q^{(1)}|^{-(qp_1/pp_1')}|Q^{(2)}|^{-(qp_1)/(pp_1')}\Big(\int\limits_{Q^{(1)}\times Q^{(2)}}|f_1(y_1^{(1)},y_1^{(2)})|(\overline{M}_{\alpha_1,\alpha_2}(v)(y_1^{(1)},y_1^{(2)}))^{1/(qm)}dy_1^{(1)}dy_1^{(2)})\Big)^{qp_1/p}\Big]^{p/p_1}$$

$$\times\cdots\times\Big[\sum_{Q^{(1)},Q^{(2)}\in\mathcal{D}}|Q^{(1)}|^{-(qp_m/pp_m')}|Q^{(2)}|^{-(qp_m/pp_m')}$$

$$\times\Big(\int\limits_{Q^{(1)}\times Q^{(2)}}|f_m(y_m^{(1)},y_m^{(2)})|(\overline{M}_{\alpha_1,\alpha_2}(v)(y_m^{(1)},y_m^{(2)}))^{1/(qm)}dy_m^{(1)}dy_m^{(2)})\Big)^{qp_m/p}\Big]^{p/p_m}$$

(by applying twice Theorem G for $w\equiv 1$ with respect to each variable separately and Minkowski's generalized inequality)

$$\le c\prod_{i=1}^{m}\Big\|\big(\overline{M}_{\alpha_1,\alpha_2}v\big)^{p/(p_iqm)}f_i\Big\|_{L^{p_i}(\mathbb{R}^{2n})}.$$

□

Remark 4.4. It is easy to see that Corollary 3.6 can be also obtained from Theorem 3.9.

Acknowledgement. The first and third authors were partially supported by the Shota Rustaveli National Science Foundation Grant (Contract Numbers D/13-23 and 31/47).

References

[1] D. R. Adams, Traces of potentials arising from translation invariant operators, *Ann. Scuola Norm. Sup. Pisa* **25**, 203–217 (1971).

[2] D. E. Edmunds, V. Kokilashvili and A. Meskhi, *Bounded and compact integral operators*, (Dordrecht). **543**. Dordrecht: Kluwer Academic Publishers (2002).

[3] J. García–Cuerva and J. L. Rubio de Francia, *Weighted norm inequalities and related topics*, North-Holland Mathematics Studies, **116** Notas de Matemtica (104). Amsterdam - New York - Oxford: North-Holland (1985).
[4] L. Grafakos, On multilinear fractional integrals, *Studia Math*. **102**, 49–56 (1992).
[5] L. Grafakos, and N. Kalton, Some remarks on multilinear maps and interpolation, *Math. Ann.* **319**, No. 1, 151–180 (2001)
[6] L. Grafakos, L. Liu, C. Perez and R. H. Torres, The multilinear strong maximal function, *J. Geom Anal.* **21**, 118–149 (2011).
[7] C. Kenig and E. Stein, Multilinear estimates and fractional integration, Math. Res. Lett. **6**, No.1, 1–15 (1999).
[8] V. M. Kokilashvili, Weighted Lizorkin–Triebel spaces. Singular integrals, multipliers, imbedding theorems, *Proc. Steklov Inst. Math.* **161**, 135-162 (1984); translation from *Tr. Mat. Inst. Steklova* **161**, 125–149 (1983).
[9] V. Kokilashvili, New aspects in weight theory, In: *Function Spaces, Differential Operators and Nonlinear Analysis. Prometeus Publishing House, Prague*, 51–70 (1996).
[10] V. Kokilashvili, M. Mastylo and A. Meskhi, On the boundedness of the multilinear fractional integral operators, *Nonlinear Anal.* **94**, 142–147 (2014).
[11] V. Kokilashvili and M. Krbec, *Weighted inequalities in Lorentz and Orlicz spaces*, Singapore etc.: World Scientific Publishing Co. Pte. Ltd. (1991).
[12] V. Kokilashvili and A. Meskhi, Two–weight estimates for strong fractional maximal functions and potentials with multiple kernels, J. Korean Math. Soc. **46**, No. 3, 523-550 (2009).
[13] V. Kokilashvili, A. Meskhi and L.-E. Persson, *Weighted norm inequalities for integral transforms with product kernels*, Mathematics Research Developments Series. New York, NY: Nova Science Publishers (2009).
[14] A. K. Lerner, S. Ombrosi, C. Pérez, R. H. Torres, R. Trujillo–González, New maximal functions and multiple weights for the multilinear Calderón-Zygmund theory, Adv. Math. **220**, No. 4, 1222–1264 (2009).
[15] K. Moen, Weighted inequalities for multilinear fractional integral operators, *Collect. Math.* **60**, 213–238 (2009).
[16] B. Muckenhoupt, Weighted norm inequalities for the Hardy maximal function, *Trans. Am. Math. Soc.* **165**, 207–226 (1972).
[17] C. Pérez, Two weighted norm inequalities for potential and fraction maximal operators, *Indiana Univ. Math. J.* **43**, 663-683 (1994).
[18] G. Pradolini, Weighted inequalities and pointwise estimates for the multilinear fractional integral and maximal operators, *J. Math. Anal. Appl.* **367**, 640–656 (2010).
[19] E. T. Sawyer, A characterization of a two-weight norm inequality for maximal operators, *Studia Math.* **75**, 1–11 (1982).
[20] E. T. Sawyer, A two-weight weak type inequality for fractional integrals. *Trans. Amer. Math. Soc.* **281**, 339-345 (1984).
[21] E. T. Sawyer and R. L. Wheeden, Weighted inequalities for fractional integrals on Euclidean and homogeneous spaces, *Amer. J. Math.* **114**, 813–874 (1992).
[22] Y. Shi and X. Tao, Weighted L^p boundedness for multilinear fractional integral on product spaces, *Analysis in Theory and Applications*, **24**, No. 3, 280–291 (2008).
[23] K. Tachizawa, On weighted dyadic Carleson's inequalities, *J. Inequal. Appl.* **6**, No. 4, 415–433 (2001).
[24] R. L. Wheeden, A characterization of some weighted norm inequalities for the fractional maximal functions, *Studia Math.* **107**, 251–272 (1993).